

TGF β 2 Polyclonal Antibody

Cat No: HR1AP4709

For research use only

Overview

Product Name	TGF β 2 Polyclonal Antibody
Source	Rabbit
Applications	WB,IHC-p,ELISA
Species Reactivity	Human,Mouse,Rat
Recommended Dilutions	
Immunogen	
Species	Rabbit
Storage	-20°C/1 year
Isotype	
Clonality	
Concentration	1 mg/ml
Observed band	44kDa
GeneID?Human?	TGFB2
Human Swiss-Prot No.	
Cellular localization	
Alternative Names	TGFB2; Transforming growth factor beta-2; TGF-beta-2; BSC-1 cell growth inhibitor; Cetermin; Glioblastoma-derived T-cell suppressor factor; G-TSF; Polyergin
Background	transforming growth factor beta 2(TGFB2) Homo sapiens This gene encodes a secreted ligand of the TGF-beta (transforming growth factor-beta) superfamily of proteins. Ligands of this family bind various TGF-beta receptors leading to recruitment and activation of SMAD family transcription factors that regulate gene expression. The encoded preproprotein is proteolytically processed to generate a latency-associated peptide (LAP) and a mature peptide, and is found in either a latent form composed of a mature peptide homodimer, a LAP homodimer, and a latent TGF-beta binding protein, or in an active form consisting solely of the mature peptide homodimer. The mature peptide may also form heterodimers with other TGF-beta family members. Disruption of the TGF-beta/SMAD pathway has been implicated in a variety of human cancers. A chromosomal translocation that includes this gene is associated with Peters's anomaly, a congenital defect of the an