

## Cleaved-Factor Xa activated HC (I235) Polyclonal Antibody

Cat No: HR1AP6300

For research use only

### Overview

Product Name	Cleaved-Factor Xa activated HC (I235) Polyclonal Antibody
Source	Rabbit
Applications	WB,ELISA
Species Reactivity	Human,Mouse,Rat
Recommended Dilutions	
Immunogen	
Species	Rabbit
Storage	-20°C/1 year
Isotype	
Clonality	
Concentration	1 mg/ml
Observed band	30kDa
GeneID?Human?	F10
Human Swiss-Prot No.	
Cellular localization	
Alternative Names	F10; Coagulation factor X; Stuart factor; Stuart-Prower factor
Background	<p>coagulation factor X(F10) Homo sapiens This gene encodes the vitamin K-dependent coagulation factor X of the blood coagulation cascade. This factor undergoes multiple processing steps before its preproprotein is converted to a mature two-chain form by the excision of the tripeptide RKR. Two chains of the factor are held together by 1 or more disulfide bonds; the light chain contains 2 EGF-like domains, while the heavy chain contains the catalytic domain which is structurally homologous to those of the other hemostatic serine proteases. The mature factor is activated by the cleavage of the activation peptide by factor IXa (in the intrinsic pathway), or by factor VIIa (in the extrinsic pathway). The activated factor then converts prothrombin to thrombin in the presence of factor Va, Ca<sup>2+</sup>, and phospholipid during blood clotting. Mutations of this gene result in factor X deficiency, a hemorrhagic condition of variable severity.</p> <p>Alternative sp</p>