

# TLE1/2/3/4 Polyclonal Antibody

Cat No: HR1AP9575

For research use only

## Overview

|                       |   |
|-----------------------|---|
| Product Name          | TLE1/2/3/4 Polyclonal Antibody  |
| Source                | Rabbit  |
| Applications          | WB,ELISA  |
| Species Reactivity    | Human,Mouse,Rat   |
| Recommended Dilutions |   |
| Immunogen             |   |
| Species               | Rabbit  |
| Storage               | -20°C/1 year  |
| Isotype               |   |
| Clonality             |   |
| Concentration         | 1 mg/ml   |
| Observed band         | 90kDa   |
| GeneID?Human?         |   |
| Human Swiss-Prot No.  |   |
| Cellular localization |   |
| Alternative Names     | similar to transducin-like enhancer of split 1/2/3/4  |
| Background            | <p>function:Transcriptional corepressor that binds to a number of transcription factors. Inhibits NF-kappa-B-regulated gene expression. Inhibits the transcriptional activation mediated by FOXA2, and by CTNNB1 and TCF family members in Wnt signaling. The effects of full-length TLE family members may be modulated by association with dominant-negative AES. Unusual function as coactivator for ESRRG.,PTM:Phosphorylated, probably by CDC2. The degree of phosphorylation varies throughout the cell cycle, and is highest at the G2/M transition. Becomes hyperphosphorylated in response to cell differentiation and interaction with HES1 or RUNX1.,similarity:Belongs to the WD repeat Groucho/TLE family.,similarity:Contains 6 WD repeats.,subcellular location:Nuclear and chromatin-associated, depending on isoforms and phosphorylation status. Hyperphosphorylation decreases the affinity for nuclear components.,subunit:Homooligomer and heterooligomer with other family members. Binds LEF1, RUNX1, RUNX3, FOXA2, KDM6A, UTY, histone H3, HESX1, ESRRG and the NF-kappa-B subunit RELA. Interacts with HES1 (via WRPW motif),tissue specificity:In all tissues examined, mostly in brain, liver and muscle.,</p> |