

DCAMKL2 Polyclonal Antibody

Cat No: HR1AP3261

For research use only

Overview

| | |
|-----------------------|---|
| Product Name | DCAMKL2 Polyclonal Antibody |
| Source | Rabbit |
| Applications | WB,IF,ELISA |
| Species Reactivity | Human,Mouse |
| Recommended Dilutions | |
| Immunogen | |
| Species | Rabbit |
| Storage | -20°C/1 year |
| Isotype | |
| Clonality | |
| Concentration | 1 mg/ml |
| Observed band | 83kDa |
| GeneID?Human? | DCLK2 |
| Human Swiss-Prot No. | |
| Cellular localization | |
| Alternative Names | DCLK2; DCAMKL2; DCDC3B; DCK2; Serine/threonine-protein kinase DCLK2; CaMK-like CREB regulatory kinase 2; CL2; CLICK-II; CLICK2; Doublecortin domain-containing protein 3B; Doublecortin-like and CAM kin |
| Background | <p>doublecortin like kinase 2(DCLK2) Homo sapiens This gene encodes a member of the protein kinase superfamily and the doublecortin family. The protein encoded by this gene contains two N-terminal doublecortin domains, which bind microtubules and regulate microtubule polymerization, a C-terminal serine/threonine protein kinase domain, which shows substantial homology to Ca²⁺/calmodulin-dependent protein kinase, and a serine/proline-rich domain in between the doublecortin and the protein kinase domains, which mediates multiple protein-protein interactions. The microtubule-polymerizing activity of the encoded protein is independent of its protein kinase activity. Mouse studies show that the DCX gene, another family member, and this gene share function in the establishment of hippocampal organization and that their absence results in a severe epileptic phenotype and lethality, as described in human patients with lissencephaly. Multiple alterna</p> |