

Met Polyclonal Antibody

Cat No: HR1AP3893

For research use only

Overview

Product Name	Met Polyclonal Antibody
Source	Rabbit
Applications	WB,IF,ELISA
Species Reactivity	Human,Mouse,Rat
Recommended Dilutions	
Immunogen	
Species	Rabbit
Storage	-20°C/1 year
Isotype	
Clonality	
Concentration	1 mg/ml
Observed band	155kDa
GeneID?Human?	MET
Human Swiss-Prot No.	
Cellular localization	
Alternative Names	MET; Hepatocyte growth factor receptor; HGF receptor; HGF/SF receptor; Proto-oncogene c-Met; Scatter factor receptor; SF receptor; Tyrosine-protein kinase Met
Background	MET proto-oncogene, receptor tyrosine kinase(MET) Homo sapiens This gene encodes a member of the receptor tyrosine kinase family of proteins and the product of the proto-oncogene MET. The encoded preproprotein is proteolytically processed to generate alpha and beta subunits that are linked via disulfide bonds to form the mature receptor. Further processing of the beta subunit results in the formation of the M10 peptide, which has been shown to reduce lung fibrosis. Binding of its ligand, hepatocyte growth factor, induces dimerization and activation of the receptor, which plays a role in cellular survival, embryogenesis, and cellular migration and invasion. Mutations in this gene are associated with papillary renal cell carcinoma, hepatocellular carcinoma, and various head and neck cancers. Amplification and overexpression of this gene are also associated with multiple human cancers. [provided by RefSeq, May 2016].