

GAK Polyclonal Antibody

Cat No: HR1AP3504

For research use only

Overview

| | |
|-----------------------|--|
| Product Name | GAK Polyclonal Antibody |
| Source | Rabbit |
| Applications | WB,IHC-p,IF,ELISA |
| Species Reactivity | Human,Mouse |
| Recommended Dilutions | |
| Immunogen | |
| Species | Rabbit |
| Storage | -20°C/1 year |
| Isotype | |
| Clonality | |
| Concentration | 1 mg/ml |
| Observed band | 144kDa |
| GeneID?Human? | GAK |
| Human Swiss-Prot No. | |
| Cellular localization | |
| Alternative Names | GAK; Cyclin-G-associated kinase |
| Background | <p>cyclin G associated kinase(GAK) Homo sapiens In all eukaryotes, the cell cycle is governed by cyclin-dependent protein kinases (CDKs), whose activities are regulated by cyclins and CDK inhibitors in a diverse array of mechanisms that involve the control of phosphorylation and dephosphorylation of Ser, Thr or Tyr residues. Cyclins are molecules that possess a consensus domain called the &#x201c;cyclin box.&#x201d; In mammalian cells, 9 cyclin species have been identified, and they are referred to as cyclins A through I. Cyclin G is a direct transcriptional target of the p53 tumor suppressor gene product and thus functions downstream of p53. GAK is an association partner of cyclin G and CDK5. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Dec 2015],</p> |